




Taletrectinib, a next-generation selective ROS1 inhibitor, inhibits growth of ROS1 wild-type and ROS1-G2032R xenografts

Hitisha K. Patel, Laura Heller, Kevin Bowman, Zheyi Hu, Yan Zhang, Irma M. Grossi, Max Pan, and Gary Hattersley

Nuvation Bio, Inc., New York, NY



Copies of this poster obtained through QR Code are for personal use only and may not be reproduced without permission from the author of this poster.

Background

- *ROS1* gene fusions occur in approximately 2% of patients with NSCLC¹
 - *ROS1* gene fusions result in increased ROS1 autophosphorylation and constitutive activation²
- While 3 ROS1 TKIs are currently approved by the FDA for the treatment of *ROS1*+ NSCLC, there remains an unmet need for effective and tolerable treatment options¹
 - Crizotinib and entrectinib are not active against many resistance mutations, including ROS1^{G2032R}, the most common mutation¹
 - Repotrectinib, while active in the CNS, is associated with a high rate of neurologic AEs such as dizziness (65%), ataxia (28%), and cognitive impairment (25%), which are attributed to the drug's inhibition of TRKB³
- Taletrectinib is a next-generation, CNS-active, ROS1 TKI with selectivity over TRKB^{4,5}
 - Taletrectinib demonstrated high and durable overall responses, robust intracranial responses, prolonged PFS, activity against G2032R, and had a favorable safety profile in the pivotal regional TRUST-I (NCT04395677)⁴ and global TRUST-II (NCT04919811)⁵ studies of *ROS1*+ NSCLC
 - NDA was accepted for priority review by the US FDA, with a PDUFA date of June 23, 2025


Methods

Study Design

- **Biochemical inhibition:** In vitro Kinase activity was detected via Reaction Biology Hotspot Kinase Assay and measured using the P81 filter-binding method⁶
- **In vitro cell viability:** Ba/F3 cells harboring the respective *ROS1* wild type or mutant fusions were plated at a density of 500–2500 cells/well and treated the next day with respective treatments. Viability was assessed after 5 or 6 days of treatment, and data are represented in IC₅₀ where 50% of growth inhibition relative to control was observed
- **Western blots:** Cells were harvested 2 hours post dosing, and protein expression was analyzed using the following antibodies: Phospho-ROS1 Tyr 2274 (CST-3028), ROS1 (CST-3287, OTI1A1-Invitrogen), and GAPDH (Proteintech, 60004). Protein expression was quantified and normalized to GAPDH expression
- **In vivo CDX/PDX:** CDX or PDX studies were run as per standard practice. Briefly, cells or tumor fragments were implanted in mice, and mice were housed in pathogen-free housing with access to sterilized food and water ad libitum. Taletrectinib, crizotinib, entrectinib, or repotrectinib was administered orally. For subcutaneous models, tumors were measured twice/week and tumor volume was calculated using the formula: (L*W²)*0.52. For intracranial models, survival of mice was evaluated


Results

Taletrectinib Demonstrates Selectivity for ROS1^{WT} and ROS1^{G2032R} Over TRKB at Physiologically Relevant ATP Concentrations




Kinase	Taletrectinib	Crizotinib	Entrectinib	Repotrectinib
ROS1 ^{WT}	1	7	4	1
ROS1 ^{G2032R}	10	2259	981	9
TRKB	92	178	1	1
	Selectivity (fold)			
TRKB/ROS1 ^{WT}	92	25	0.25	1
TRKB/ROS1 ^{G2032R}	9	<0.1	<0.001	<0.1

Taletrectinib Inhibits Cell Growth in Models Harboring ROS1^{WT} and ROS1^{G2032R} Fusions




	Cell Line	Taletrectinib	Crizotinib	Entrectinib	Repotrectinib
CD74	ROS1 ^{WT}	11	63	28	10
	ROS1 ^{G2032R}	67	1198	1695	35
SLC34A2	ROS1 ^{WT}	10	31	25	4
	ROS1 ^{G2032R}	35	620	1428	12

Taletrectinib Demonstrates Growth Inhibition and Tumor Regression in Models Harboring ROS1^{WT} and ROS1^{G2032R} Fusions




Taletrectinib Demonstrates Activity Against Single and Double ROS1^{G2032R} Mutants Resistant to Crizotinib and Entrectinib




	SLC34A2-ROS1	SLC34A2-ROS1-G2032R	SLC34A2-ROS1-G2032R-D2113G	SLC34A2-ROS1-G2032R-M2134I	SLC34A2-ROS1-G2032R-F2075V	SLC34A2-ROS1-L2026M	SLC34A2-ROS1-S1969F	SLC34A2-ROS1-D2113G	SLC34A2-ROS1-D2113N	SLC34A2-ROS1-E2020K
Taletrectinib	10	33	65	8	23	31	1	3	3	10
Crizotinib	32	721	509	940	548	107	6	10	6	76
Entrectinib	26	1419	2132	903	2947	63	2	10	3	29
Repotrectinib	5	12	19	4	10	12	<1	1	1	5

Taletrectinib Prolongs Survival in LU-01-0414 Intracranial Mouse PDX Model Harboring a SDC4-ROS1 Fusion



Taletrectinib Inhibits ROS1 Phosphorylation in Both ROS1^{WT} and ROS1^{G2032R} Resistance Mutation Settings



Conclusions

- Taletrectinib demonstrates selectivity for ROS1^{WT} and ROS1^{G2032R} over TRKB at physiologically relevant ATP concentrations, whereas repotrectinib, crizotinib, and entrectinib did not demonstrate any selectivity over TRKB, a kinase whose inhibition is associated with CNS toxicity
- Taletrectinib demonstrated in vitro and in vivo activity against ROS1 regardless of fusion partner or resistance mutation, including the ROS1^{G2032R} mutation which is resistant to crizotinib and entrectinib
- Taletrectinib demonstrated tumor regression in models harboring ROS1^{WT} and ROS1^{G2032R} fusions
- Taletrectinib inhibits ROS1 phosphorylation in ROS1^{WT} and ROS1^{G2032R} models
- Taletrectinib treatment prolonged survival in mice implanted intracranially with a PDX model harboring a SDC4-ROS1 fusion

Acknowledgments

- This study was sponsored by Nuvation Bio, Inc. Writing and graphical support were provided by David M. Jensen, PhD, of Peloton Advantage, LLC, an OPEN Health company, and funded by Nuvation Bio.

References

1. Boulanger MC, et al. *Oncologist*. 2024;29:943-956.
2. Davies KD, et al. *Clin Canc Res*. 2013;19:4040-4045.
3. Augtyro [package insert]. Princeton, NJ: Turning Point Therapeutics, Inc., a Bristol Myers Squibb company; 2024.

4. Li W, et al. *J Clin Oncol*. 2024;42:2660-2670.
5. Perol M, et al. *Ann Oncol*. 2023;34(suppl 2):S788-S789.
6. Anastassiadis T, et al. *Net Biotechnol*. 2011;29:1039-1045.
7. Hui Q, et al. *Cancer Res*. 2022;82(suppl 12):5989.