

Taletrectinib vs Entrectinib in *ROS1*-Positive Non-Small Cell Lung Cancer: A Matching-Adjusted Indirect Comparison

Misako Nagasaka,¹ Geoffrey Liu,² Nathan A. Pennell,³ Maurice Pérol,⁴ Wenfeng Chen,⁵ Lyudmila Bazhenova,⁶ Caicun Zhou⁷

¹University of California Irvine School of Medicine and Chao Family Comprehensive Cancer Center, Orange, CA, USA; ²Princess Margaret Cancer Centre, Temerty School of Medicine, University of Toronto, Toronto, Canada; ³Taussig Cancer Institute, Cleveland Clinic Cancer Center, Cleveland, OH, USA; ⁴Centre Léon Bérard, Lyon, France; ⁵Nuvation Bio, New York, NY, USA; ⁶University of California San Diego Moores Cancer Center, San Diego, CA, USA; ⁷Shanghai Pulmonary Hospital and Thoracic Cancer Institute, Tongji University School of Medicine, Shanghai, China



Copies of this poster obtained through QR code are for personal use only and may not be reproduced without permission from the author of this poster.

Objective

To conduct a matching-adjusted indirect comparison (MAIC) of taletrectinib and entrectinib in TKI-naïve patients with *ROS1*+ NSCLC

Background

- For patients with *ROS1*+ NSCLC, *ROS1* tyrosine kinase inhibitors (TKIs) are the current standard of care¹
- Entrectinib was approved by the US Food and Drug Administration in 2019 and crosses the blood-brain barrier, but overall responses were marginal in patients with prior CNS progression (IC-ORR: 11%)^{2,3}
- Taletrectinib is an oral, potent, CNS-active, selective, next-generation, *ROS1* inhibitor⁴⁻⁶
 - Taletrectinib has demonstrated high and durable overall and IC response rates, activity against G2032R, and a favorable safety profile in the pivotal regional TRUST-I (NCT04395677) and global TRUST-II (NCT04919811) studies^{6,7}
 - In December 2024, the US FDA granted priority review to the NDA for seeking the approval of taletrectinib for the treatment of advanced *ROS1*+ NSCLC (PDUFA goal date of June 23, 2025)
- In the absence of head-to-head trials in TKI-naïve patients with *ROS1*+ NSCLC, we compared taletrectinib with the first-generation TKI, entrectinib, using a MAIC analysis

Abbreviations

CNS, central nervous system; CR, complete response; DOR, duration of response; ECOG PS, Eastern Cooperative Oncology Group performance status; IC, intracranial; MAIC, matching-adjusted indirect comparison; NA, not available; NDA, New Drug Application; NSCLC, non-small cell lung cancer; ORR, objective response rate; OS, overall survival; PD, progressive disease; PDUFA, Prescription Drug User Fee Act; PFS, progression-free survival; PR, partial response; PT, preferred term; *ROS1*+, *ROS1*-positive; SAE, serious adverse events; SD, stable disease; TKI, tyrosine kinase inhibitor; TRAE, treatment-related adverse event; US FDA, US Food and Drug Administration.

References


1. Gendarme S, et al. *Curr Oncol*. 2022;29:641-658. 2. Rozlytrek [package insert]. San Francisco, CA: Genentech USA, Inc; October 2023. 3. Drilon A, et al. *JTO Clin Res Rep*. 2022;3:100332. 4. Nagasaka M, et al. *Future Oncol*. 2023;19:123-135. 5. Katayama R, et al. *Nat Comm*. 2019;10:3604. 6. Li W, et al. *J Clin Oncol*. 2024;42:2660-2670. 7. Liu, et al. *JTO*. 2024;10:S72-S73. 8. Drilon A, et al. *Lancet Oncol*. 2020;21:261-270. 9. Fan Y, et al. *Clin Lung Cancer*. 2024;25:e81-e86.e4. 10. Perol M, et al. *JCO*. 2025.

Acknowledgments

- We would like to thank all patients who participated in this study, the study investigators, and their staff
- This study was sponsored by Nuvation Bio, Inc. Writing and graphical support were provided by Tulika Bhushan Bahukhandi, RPh, MS, of Peloton Advantage, LLC, an OPEN Health company, and funded by Nuvation Bio


Methods

Study Design




Results


Baseline Comparison in *ROS1*+ TKI-Naïve NSCLC




ORR in *ROS1*+ TKI-Naïve NSCLC




OS for *ROS1*+ TKI-Naïve Group



PFS for *ROS1*+ TKI-Naïve Group



DOR for *ROS1*+ TKI-Naïve Group



Safety in *ROS1*+ NSCLC

	Taletrectinib (N=352)*	Entrectinib ⁹ (N=247)
TRAEs	n (%)	n(%)
TRAEs leading to permanent discontinuation	9 (2.6)	17 (6.9)
TRAEs leading to treatment interruption	96 (27.3)	89 (36.0)
TRAEs leading to dose modification	97 (27.6)	86 (34.8)
Any grade 3/4 TRAEs	116 (33.0)	107 (43.3)
Treatment-related SAEs	27 (7.7)	35 (14.2)
Treatment-related deaths	3 (0.9)	1 (0.4)

* Pooled safety population included all patients receiving taletrectinib 600 mg once daily until disease progression or unacceptable toxicity across the safety population¹⁰

- TRAЕ rates are summarized for taletrectinib and entrectinib
- Due to the inconsistent definition of TRAЕs, as well as lack of availability of baseline patient characteristics of comparable safety population, these estimates are unadjusted and not designed for drawing comparisons between agents

Limitations

- Indirect treatment comparison studies are subject to bias due to variations in patient populations, study designs, and outcome measures across the included trials, leading to uncertain comparisons

Strength

- Given the available evidence, the described comparative analysis is a transparent and methodologically sound approach to compare taletrectinib and entrectinib

Conclusions

Taletrectinib showed significantly improved efficacy outcomes vs entrectinib in TKI-naïve patients with *ROS1*+ NSCLC in the MAIC analysis, including

- Higher ORR and a significant 65% improvement in the likelihood of maintaining a response
- Significant 52% reduction in the risk of death and a 58% reduction in the risk of progression/death

TRAЕs are summarized as unadjusted estimates, since the lack of comparable baseline characteristics limits comparison across treatments

These findings underscore the therapeutic benefits of taletrectinib over entrectinib outside of head-to-head randomized controlled trials